

This article was given to us by the [Tea Council](#).

## **Tea and Cardiovascular Disease**

### ***Introduction***

Cardiovascular Disease (CVD) is the main cause of death in the UK, accounting for over 250,000 deaths a year, more than one in three people. The main forms of CVD are Coronary Heart Disease (CHD) and stroke. About half of all deaths from CVD are from CHD and about a quarter are from stroke. More than 110,000 people die of heart problems, 300,000 have heart attacks and 1.4 million suffer from angina every year. In addition, CHD accounts for about 3% of all hospital admissions in England.

### ***Risk Factors associated with CHD***

There is no one single cause for CHD, instead the risk of developing it is determined by a combination of factors, some of which are unalterable, others that are potentially modifiable. These risk factors are outlined below.

### **Modifiable Risk Factors--Non-modifiable risk factors**

Hyperlipidaemia (raised cholesterol and triglycerides)

Hypertension (high blood pressure)

Reduced physical activity

Smoking

Stress

Obesity

Family history

Sex

Age

Other physiological factors and conditions that may increase the risk of developing CHD include:-

Insulin resistance syndrome

Oxidation of lipoproteins

Inflammation and endothelial damage

Hyperhomocysteinaemia (high blood homocysteine)

Platelet aggregation

Clotting factors

Diabetes Mellitus

Renal Disease

### ***Prevention***

Primary prevention of CVD involves adopting a 'healthy lifestyle' to control the 'Modifiable' risk factors. The important aspects of this type of lifestyle include:-

- ~ Quitting smoking
- ~ Taking regular exercise
- ~ Adopting a healthy diet
- ~ Reducing stress and anxiety levels

### ***Dietary Recommendations***

- ~ A healthy diet that includes the following;-
- ~ Increasing the consumption of oily fish
- ~ Increasing fruit, vegetable and pulses
- ~ Replacing some of the dietary energy coming from total fat with complex starchy foods
- ~ Moderating fat reduction with the use of monounsaturated fats
- ~ Limiting salt intake
- ~ Consuming modest amounts of alcohol can help to reduce the risk of CVD by altering a range of risk factors in a number of ways. Some of these effects are outlined below.

### **Dietary Intervention--Physiological Effect**

- ~ Omega 3 fatty acids (oily fish, rapeseed oil)
- ~ Anti-inflammatory, protecting against endothelial damage
- ~ Anti-thrombotic
- ~ Reduces triglycerides
- ~ Improves insulin sensitivity
- ~ Anti-arrhythmic effects
- ~ Other fatty acids Action of omega-3 fatty acids is improved by lower intakes of saturated fat
- ~ Lower saturated fats reduces the risk of thrombosis and reduces serum lipids
- ~ Fruit, vegetables, pulses, nuts
- ~ Antioxidants protect against Low Density Lipoprotein (LDL) oxidation
- ~ Rich in potassium may help to control blood pressure
- ~ Rich in folic acid, may prevent homocysteine formation
- ~ Contain soluble fibre, helping to improve lipid profiles

### **Alcohol**

- ~ Improved HDL levels
- ~ Possible antioxidant effects

### ***The benefits of tea***

There is growing evidence suggesting that other dietary components may enhance the general healthy eating recommendations. For example, research is highlighting that there maybe a beneficial relationship between tea drinking and reduced risk of cardiovascular disease.

### ***The evidence for tea and CVD***

A number of epidemiological studies have demonstrated that total flavonol/flavonoid or tea consumption is linked to protection from heart disease and stroke. However, not all

studies have shown similar protective effects. Inconsistencies from these studies may be as a result of their design e.g. lack of detail about exposure to tea – quantity, strength and variety, insufficient information about the flavonoid contents of foods and other risk factors for CVD not being considered. These details may influence the end results and consideration of these factors is required for any future research.

However, the conclusions from a recent meta-analysis of all studies investigating the relationship between tea drinking and CHD suggested that drinking 3 cups of tea a day reduces the risk of myocardial infarction by 11%.

### ***Protective mechanisms of tea***

The outcomes from these studies have created great interest in understanding the mechanisms by which tea may confer its cardiovascular protective properties.

Mechanisms such as the prevention of LDL oxidation, reduction in blood lipids, anti-inflammatory actions, improvements in blood vessel function and inhibition of platelet aggregation have been established mainly in animal and in vitro studies. These mechanisms are discussed in greater detail in the fact sheet 'Tea and 'Antioxidants'. Results in human studies are not as clear, and although some epidemiological studies seem to indicate that tea can reduce serum cholesterol, most studies show no effect in reducing serum cholesterol or blood pressure. As antioxidant activity from tea has been demonstrated in vivo, it may be that tea exerts its cardioprotective properties in other ways such as improvements in endothelial dysfunction, platelet aggregation or ways that have yet to be investigated.

### ***In summary***

From a number of epidemiological studies investigating the relationship between tea drinking and cardiovascular disease, it is reasonable to conclude that drinking both green and black tea is compatible with dietary advice to help protect against CVD. However, although the scientific evidence for the mechanisms by which tea exerts its positive health effects is growing, it is not yet conclusive and represents promising areas for future research in human trials.

### **References:**

1. British Heart Foundation. Coronary Heart Disease Statistics Database, 2000
2. National Service Frameworks Coronary Heart Disease, Department of Health, 2000
3. Department of Health (1994) Nutritional Aspects of Cardiovascular Disease, Report of the Cardiovascular Review Group of the Committee of Medical Aspects of Food Policy (COMA). Report No. 46. London: HMSO
4. British Nutrition Foundation. Diet and Heart Disease: A round table of factors. London: Chapman and Hall, 1997
5. Sesso HD, et al. (1999) Coffee and tea intake and the risk of myocardial infarction. *Am J Epidemiol*, Jan 15, 149(2), 162-167
6. Keli SO, et al. (1996) Dietary flavonoids, antioxidant vitamins, and incidence of stroke: the Zutphen study. *Arch Intern Med*, 156(6), 637-642

7. Knekt P, et al. Flavonoid intake and coronary mortality in Finland: a cohort study. *BMJ*, 312, 478-81, 1996
8. Stensvold I, et al. Tea consumption. Relationship to cholesterol, blood pressure and coronary and total mortality. *Preventative Med*, 21, 546,1992.
9. Yochum L, et al. Dietary flavonoid intake and risk of cardiovascular disease in postmenopausal women. *Am J Epidemiol*, 149(10), 943-9, 1999
10. Hertog MGL, et al. (1993) Dietary antioxidants flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *Lancet*, 155,381
11. Hertog MGL, et al. (1995) Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. Hertog, et al, *Arch Int Med* 155,381
12. Geleijnse JM, et al (1999) Tea flavonoids may protect against atherosclerosis: the Rotterdam study. *Arch Intern Med* 159; 2170-4
13. Hertog MG, et al (1997) Antioxidant flavonols and ischaemic heart disease in a Welsh population of men: the Caerphilly study. *Am J Clin Nutr*; 65: 1489-94
14. Rimm EB, et al (1996) Relation between intake of flavonoids and risk for coronary heart disease in male health professionals. *Ann Intern Med*; 125: 384-9
15. Woodward M, et al (1999) Coffee and tea consumption in the Scottish Heart Health Study follow up: conflicting relations with coronary risk factors, coronary disease and all cause mortality. *J Epidemiol Comm Health*: 53: 481-7
16. Klatsky AL, et al (1990) Coffee use prior to myocardial infarction restudied: heavier intake may increase the risk. *Am J Epidemiol* 132; 479-88
17. Peters U, et al (2001) Does tea affect Cardiovascular Disease? A meta-Analysis. *Am J Epidemiol* 154; 6: 495-501
18. Vinson JA, et al (1995) Plant flavonoids, especially tea flavonols, are powerful antioxidants using an in vitro oxidation model for heart disease. *J Agric Food Chem*; 43 (11):2800-2802
19. De Whalley, et al (1990) Flavonoids inhibit the oxidative modification of Low Density Lipoproteins by macrophages. *Biochem Pharmacol* 39; 1743-1750
20. Yoshida H, et al (1999) Inhibitory effect of tea flavonoids on the ability of cells to oxidise low density lipoprotein. *Biochem Pharmacol* 58; 1695-703 *Biochem Pharmacol* 39; 1743-1750
21. Pearson DA, et al (1998) Inhibition of endothelial cell mediated low density lipoprotein oxidation by green tea extracts. *J Agric Food Chem* 46; 1445-9
22. Zhu QY, et al (2000) Interaction between flavonoids and alpha-tocopherol in human low density lipoprotein. *J Nutr Biochem* 11; 14-21
23. Osada K, et al (2001) Tea catechins inhibit cholesterol oxidation accompanying oxidation of low density lipoprotein in vitro. *Comp Biochem Physiol C Toxicol Pharmacol*, 128(2): 153-64
24. Yang TTC, et al (1997) Hypocholesterolemic effects of Chinese tea. *Pharmacol res* 35; 505-12
25. Lin YL, et al (1998) Hypolipidemic effect of green tea leaves through induction of antioxidant and Phase II enzymes including superoxidase dismutase, catalase, and glutathione S-transferase in rats. *J Agric food Chem* 46; 1893-9
26. Vinson JA, et al (1998) Effect of green and black tea supplementation on lipids, lipid oxidation and fibrinogen in hamster: mechanisms for the epidemiological benefits of tea drinking. *FEBS Lett* 433; 44-6

27. Tijburg LBM (1997) Tea flavonoids and cardiovascular diseases a review. *Crit Rev Food Sci. Nutr* 37; 771-85
28. Hofbauer R, et al (1999) The green tea extract epigallocatechin is able to reduce neutrophil transmigration through monolayers of endothelial cells. *Wiener Klinische Wochenschrift* 111; 278-82
29. Duffy SJ, et al. (2001) Short and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation*, 10, 104, 2, 151
30. Corvazier E, et al (1985) Interference of some flavonoids and non-steroidal anti-inflammatory drugs with oxidative metabolism of arachidonic acid by human platelets and neutrophils. *Biochemica et Biophysica Acta* 835; 315-321
31. Polette A, et al (1996) N-3 fatty acid-induced lipid peroxidation in human platelets is prevented by catechins. *Thrombosis and Haemostasis* 95; 945-9
32. Kelly C, et al (1996) Modulation of human platelet function by food flavonoids. *Biochemical Soc transactions* 24; 197
33. Tzeng SH, et al (1991) Inhibition of platelet aggregation by some flavonoids. *Thromb Res*; 64: 91-100
34. Hodgson JM, et al (2001) Effects of regular ingestion of black tea on haemostasis and cell adhesion molecules in humans. *Eur J Clin Nutr*; 55 (10): 881-6
35. Kono S, et al (1996) Relation of green tea consumption to serum lipids and lipoproteins in Japanese men. *J of Epidemiology*; 6(3): 128-33
36. Kono S, et al (1992) Green tea consumption and serum lipid profiles: a cross-sectional study in northern Kyushu, Japan. *Prev Med*, 21(4): 526-31
37. Imai K, et al (1995) Cross sectional study of effects of drinking green tea on cardiovascular and liver diseases. *BMJ* 310; 693-6
38. Serafini M, et al (1996) In vivo antioxidant effect of green and black tea in man. *Eur J Clin Nutr*; 50: 28-32
39. Benzie FF, et al (1999) Consumption of green tea causes rapid increase in plasma antioxidant power in humans. *Nutr and Cancer*; 34(1): 83-87